

PHY201 List of eExam Questions in the Bank

Latex formatted questions may not properly render

Q1 A vector is completely defined by quantities, each with its appropriate units

Q2 $\frac{\partial H}{\partial p_k} = \dot{q}_k$, $\frac{\partial H}{\partial q_k} = -\dot{p}_k$ and $\frac{\partial H}{\partial t} = -\frac{\partial L}{\partial t}$ are known as the equations of Hamilton

Q3 The equation $H = \sum_{k=1}^n p_k \dot{q}_k - L(q_k, \dot{q}_k, t)$ gives the classical Hamiltonian function for a system of particles, each of mass m_k and described by the generalized coordinates q_k . For a system of particles in a conservative field, the Hamiltonian function given represents of the system

Q4 $\frac{d}{dt} \left(\frac{\partial L}{\partial \dot{q}_k} \right) = \frac{\partial L}{\partial q_k}$ represent the 's equations of motion for a conservative system subject to, at worst, only holonomic constraints

Q5 In the application of the variational principle to a physical system, the quantity L in the equation $L = T - V$ is called the

Q6 Given the $Q_k = F_i \frac{\partial x_i}{\partial q_k}$, Q_k represents

Q7 $p_k = \frac{\partial T}{\partial \dot{q}_k}$ represents momentum

Q8 Restrictions imposed on the free motion of a particle (or a system of particles) are generally called

Q9 The degree of is important in the design of a good suspension system of a car to ensure comfortable ride.

Q10 In a series LC circuit, the quantity $\frac{1}{LC}$ is equal to the square of the (where L and C are the inductance and capacitance respectively)

Q11 The equation of a simple pendulum is given as $m\ddot{x} + mg\frac{x}{l} = 0$. The quantity $\frac{g}{l}$ is equal to the of the

Q12 If the displacement of a simple harmonic oscillator as a function of time t given as $x = A \cos(\omega t + \phi)$, the quantity ϕ is called the constant

Q13 $\Delta \phi = \int \mathbf{F} \cdot d\mathbf{r}$. Note ϕ is the potential in a conservative force field.

Q14 for small displacements from equilibrium position, the restoring force of a simple harmonic oscillator obeys law

Q15 (1) The paths of planets about the sun are elliptical in shape, with the center of the sun being located at one focus. (2) An imaginary line drawn from the center of the sun to the center of the planet will sweep out equal areas in equal intervals of time. (3) The ratio of the squares of the periods of any two planets is equal to the ratio of the cubes of their average distances from the sun. The three statements constitute what is known as 's laws

Q16 The motion of a particle in a central force field always takes place in a

Q17 In the equation $\vec{N} = \vec{r} \times \vec{F}$, $\vec{N}$ represents

Q18 For a particle moving under the central conservative force, the equation $\frac{1}{2} \dot{r}^2 + \frac{L^2}{2mr^2} + V(r) = E$ is called the energy equation

Q19 In the equation $\vec{L} = \vec{r} \times \vec{p}$, $\vec{L}$ represents

Q20 The equations
$$\dot{x}_1 = \sum_{k=1}^{3N} \frac{\partial x_1}{\partial q_k} \dot{q}_k$$
$$\dot{x}_N = \sum_{k=1}^{3N} \frac{\partial x_N}{\partial q_k} \dot{q}_k$$
 represent the cartesian velocity components expressed in terms of general

Q21 If a system of particles is described by a set of generalised coordinates $q_1, \dots, q_{3N}$, then $\dot{q}_k$ for $k=1, 2, \dots, 3N$ is called the generalized associated with the coordinates

Q22 The generalised force associated with angular quantities is a

Q23 A particle that moves along a straight line has degree of freedom

Q24 Constraints that can be represented as functions of space and time are said to be

Q25 The mathematical term used for describing the circulation of a vector is the

Q26 If $\vec{a} \times \vec{b} = 0$, then, $\vec{a}$ and $\vec{b}$ are

Q27 The measure of the outward flow of the vector from its source is the of the vector

Q28 The dot or scalar product of a force and a displacement vectors defines

Q29 A force which acts on a particle in such a way that there is interconversion between kinetic and potential energies is said to be

Q30 If the addition of vector by geometric method gives a close figure, the the

resultant of the vectors is

Q31 If $\vec{F}$ is a force and $\vec{r}$ the radius vector from the origin of the position of a particle acted upon the force, then $\vec{F} \times \vec{r}$ represents the _____ of the force about the origin

Q32 The gradient of a scalar field function is a _____ quantity

Q33 If the curl of a vector is zero, then the vector is said to be _____

Q34 Undisturbed orbital motion under the influence of a central force satisfies _____'s law of areas

Q35 The term $r \frac{d^2\theta}{dt^2}$ in the expression $\frac{d^2r}{dt^2} - r \frac{d^2\theta}{dt^2}$ represents _____ acceleration

Q36 The aspect of mechanics that describes the motion of particles without that regard to the dynamical laws that determine which motion actually occurs is referred to as _____

Q37 If $\oint \vec{F} \cdot d\vec{r} = 0$, then $\vec{F}$ is a _____ force

Q38 The statement $T + V = E(\text{constant})$, where T is the kinetic energy and V the potential energy is known as the _____ theorem

Q39 If $\nabla \times \vec{F} = 0$, then $\vec{F}$ is _____

Q40 If the work done in moving a particle from place to place in a force field is independent of the path followed by the particle, then the force field is said to be _____

Q41 A damped oscillation driven by external periodic impulses whose frequency is the same as the natural frequency of the oscillator is called _____

Q42 If the resistance to a simple harmonic oscillation is very strong, the motion is said to be heavily _____

Q43 In reality, the amplitude of a simple harmonic oscillator gradually decreases in the presence of dissipative forces. Such motion is said to be _____

Q44 In an LC circuit, the quantity $\frac{1}{\sqrt{LC}}$ represents _____

Q45 The displacement of a simple harmonic oscillator is given as $x = A \cos(\omega t + \phi)$. The quantity ω is the _____ of the oscillation. (symbols have their usual meaning)

Q46 The term normal mode frequency is associated with _____ oscillation

Q47 The inertial factor in the spring mass system is m . The inertial factor in the LC circuit is

Q48 Any motion that repeats itself at regular intervals of time is said to be

Q49 The equation $\left(\frac{du}{d\theta}\right)^2 + U^2 = \frac{2m}{L^2} \left(E - V\right)$ is the statement of conservation of

Q50 For an oscillating system, if the restoring force obeys Hooke's law for small displacement from equilibrium position, the motion is

Q51 A damped oscillation driven by external periodic impulses whose frequency is the same as the natural frequency of the oscillator is said to -----

be heavily damped

free

resonate

be critically damped

Q52 The equation of motion of a simple harmonic oscillator is $\frac{d^2x}{dt^2} + 16x = 0$. Solve this equation subject to the initial conditions that at $t=0$, $x=4$ cm and $\frac{dx}{dt}=0$.

$x(t) = 3\cos 2t$

$x(t) = 4\cos 4t$

$x(t) = 2\cos t$

$x(t) = \sqrt{2}\cos\left(2t - \frac{\pi}{3}\right)$

Q53 Which of the following is NOT a valid solution of the equation of a simple harmonic oscillator? The symbols have their usual meaning.

$x = \cos(\omega t + \phi)$

$x = \sin(\omega t - \phi)$

$x = \cos(\omega t - \phi)$

$x = \sin(\omega t + \phi)$

Q54 The equation of a simple harmonic motion is given as $\frac{d^2x}{dt^2} + \omega^2 x = 0$ where the symbols have their usual meaning. The dimension of the quantity ω^2 is

L^{-1}

M

T^{-2}

LT^{-2}

Q55 Write down the correct relation for the component of the acceleration in the $\hat{r}$ direction in the plane polar coordinate system

$r\ddot{\theta} + 2\dot{r}\dot{\theta}$

$\ddot{r} + r\dot{\theta}^2$

$r\ddot{\theta} - 2\dot{r}\dot{\theta}$

$\ddot{r} - r\dot{\theta}^2$

Q56 In the plane polar coordinate system, which of the following correctly

gives the component of acceleration in the $\hat{\theta}$ direction?

$$\begin{aligned} & r\ddot{\theta} + 2\dot{r}\dot{\theta} \\ & \ddot{r} + r\ddot{\theta}^2 \\ & r\ddot{\theta} - 2\dot{r}\dot{\theta} \\ & \ddot{r} - r\ddot{\theta}^2 \end{aligned}$$

Q57 In the plane polar coordinate system, which of the following correctly gives the component of velocity in the $\hat{\theta}$ direction?

$$\begin{aligned} & \dot{r} \\ & \dot{\theta} \\ & r\dot{\theta} \\ & r\dot{r} \end{aligned}$$

Q58 if $\hat{r}$, $\hat{\theta}$ are the functions of θ in the plane polar coordinate system, which of the following is Not a correct relation among θ , $\hat{r}$, and $\hat{\theta}$?

$$\begin{aligned} & \hat{r} = \hat{x}\cos\theta - \hat{y}\sin\theta \\ & \hat{r} = \hat{x}\cos\theta + \hat{y}\sin\theta \\ & \hat{\theta} = -\hat{x}\sin\theta + \hat{y}\cos\theta \\ & \hat{\theta} = \hat{y}\cos\theta - \hat{x}\sin\theta \end{aligned}$$

Q59 In the plane polar coordinates, which of the following relations is NOT correct?

$$\begin{aligned} & x = r\cos\theta \\ & \theta = \sin^{-1} \frac{y}{\sqrt{x^2 + y^2}} \\ & \theta = \cos^{-1} \frac{x}{\sqrt{x^2 + y^2}} \\ & \theta = r\sin x \end{aligned}$$

Q60 A particle is subjected to a force $F = (2xy + z^2)\hat{i} + x\hat{j} + 2xz\hat{k}$ in a conservative field. Find the potential function.

$$\begin{aligned} & 2xyz \\ & -\left(x^2y + z^2x\right) \\ & 2x^2yz^2 \\ & -\left(xy^2 + zx^2\right) \end{aligned}$$

Q61 Given that the hamiltonian function for the motion of a one-dimensional harmonic oscillator is $H = \frac{1}{2m}p^2 + \frac{k}{2}x^2$, which of the following is the correct set of hamiltonian equation of motion for the system?

$$\begin{aligned} & \frac{dp}{m} = \dot{x}, \quad kx = -\dot{p} \\ & \frac{p}{m} = \dot{x}, \quad kx = -\dot{p} \\ & \frac{x}{m} = \dot{p}, \quad kp = -\dot{x} \\ & \frac{x}{m} = \dot{p}, \quad kx = -\dot{x} \end{aligned}$$

Q62 Which of the following is the correct Hamiltonian function for the motion of a particle in a central field?

$$\begin{aligned} & H = \frac{1}{2} \left(p_\theta^2 + \frac{p_r^2}{r^2} \right) + V(\theta) \\ & H = \frac{1}{2} \left(p_\theta^2 + \frac{p_r^2}{r^2} \right) + V(r) \\ & H = \frac{1}{2} \left(p_\theta^2 - \frac{p_r^2}{r^2} \right) + V(r) \end{aligned}$$

\$\$

$$H = \frac{1}{2} \left(p^2_r + \frac{p^2_{\theta}}{r^2} \right) + V(r)$$

Q63 Which of these is the correct set up of the Lagrangian for a simple pendulum?

$$L(\theta, \dot{\theta}) = \frac{1}{2} m l^2 \dot{\theta}^2 + m g l \cos \theta$$

$$L(\theta, \dot{\theta}) = \frac{1}{2} m l^2 \dot{\theta}^2 + m g l \sin \theta$$

$$L(\theta, \dot{\theta}) = \frac{1}{2} m l^2 \dot{\theta}^2 + m g l \cos \theta$$

$$L(\dot{\theta}) = \frac{1}{2} m l^2 \dot{\theta}^2 - m g l \cos \theta$$

Q64 Which of the following is NOT a correct canonical equation of Hamilton? (Symbols have their usual meaning)

$$\frac{\partial H}{\partial p_{\alpha}} = \dot{p}_{\alpha}$$

$$\frac{\partial H}{\partial p_{\alpha}} = \dot{q}_{\alpha}$$

$$\frac{\partial H}{\partial q_{\alpha}} = -\dot{p}_{\alpha}$$

$$\frac{\partial H}{\partial t} = -\frac{\partial L}{\partial t}$$

Q65 The equations

$$\dot{x}_1 = \sum_{k=1}^{3N} \frac{\partial x_1}{\partial q_k} \dot{q}_k$$

$$\dot{x}_N = \sum_{k=1}^{3N} \frac{\partial x_N}{\partial q_k} \dot{q}_k$$

expressed in terms of

generalised coordinates and velocities

generalised forces and acceleration

generalised forces and torques

generalised forces and momenta

Q66 If a system of particles is described by a set of generalised coordinates

$q_1, \dots, q_{3N}$, then $\dot{q}_k$ for $k=1, 2, \dots, 3N$ is called the -----

- associated with the coordinates

specific torque

generalised velocity

angular acceleration

generalised force

Q67 Which of these is an example of a system of particles subject to constraints?

a single particle

3N free particles

rigid object

N free particles

Q68 The generalised force associated with angular quantities is

potential

energy

power

a torque

Q69 In spherical coordinates, which of the following relations is NOT correct?

$$x = r \cos \theta \cos \phi$$

$$x = r \sin \theta \cos \phi$$

$$y = r \sin \theta \sin \phi$$

$$z = r \cos \theta$$

Q70 Two masses m_1 and m_2 connected by a spring are sliding on a frictionless plane. How many degrees of freedom has the system?

2

4

1

3

Q71 With polar coordinates (r, ϕ) as generalised coordinates, the generalised velocities are $(\dot{r}, \dot{\phi})$, while the velocity vector is -----

$$\dot{r} \hat{r} + r \dot{\phi} \hat{\phi}$$

$$\dot{r} \hat{r} + r \dot{\phi} \hat{r}$$

$$\dot{r} \hat{r} - r \dot{\phi} \hat{\phi}$$

$$\dot{r} \hat{r} - r \dot{\phi} \hat{r}$$

Q72 How many degrees of freedom has a particle that moves along a straight line?

3

2

1

0

Q73 Constraints that can be represented as functions of space and time are said to be -----

holonomic

orthogonal

inertial

invariant

Q74 A set of coordinate axes with respect to which the position and orientation of an object can be measured is called the -----

constraint

degree of freedom

orthogonality

frame of reference

Q75 Consider two capacitively coupled circuits shown in the figure. How many normal modes are we supposed to have in the system?

1

2

3

0

Q76 The equation of motion of a simple pendulum is given as $\ddot{\theta} + \omega^2 \theta = 0$. The equation of motion of a simple pendulum is given as $\ddot{\theta} + \omega^2 \theta = 0$. $\omega^2 =$ -----

g

$\sqrt{g/l}$

$\frac{g}{l}$

$\sqrt{\frac{l}{g}}$

Q77 The differential equation for the orbit of motion in an inverse square law force field is

$\frac{du}{d\theta} + u = \frac{k}{h^2}$

$\frac{d^2 u}{d\theta^2} + u = \frac{k}{h}$

$\frac{d^2 u}{d\theta^2} + u = 0$

$\frac{d^2 u}{d\theta^2} + u = \frac{k}{h^2}$

Q78 The equation of motion of a damped oscillator is given as $m\ddot{x} + \gamma\dot{x} + kx = 0$. γ is called the damping coefficient. Which of the following is the correct unit of the damping coefficient?

kg s^{-1}

ms^{-1}

kg m^{-1}

s^{-2}

Q79 All bodies at the same place in a gravitational field experience

same acceleration

different acceleration

zero acceleration

twice the acceleration as that on the surface of the source of the field

Q80 Which of the following is a correct example of a coupled oscillator?

an oscillating torsional pendulum

oscillating atoms in a solid

an acoustic oscillator

an oscillating simple pendulum

Q81 Given that $\vec{F} = f(r)\vec{r}$, where $\vec{F}$ is the central force, $f(r)\vec{r}$ is a function of the radial distance from the origin and $\vec{r}$ is the unit vector directed radially outward from the origin, which of the following is correct?

$f(r) < 0$ implies that the $\vec{F}$ is repulsive

$f(r) > 0$ implies that the $\vec{F}$ is attractive

$\vec{r} \cdot \vec{F} < 0$

$\vec{r} \cdot \vec{F} = 0$

Q82 Which of the following is NOT a central force?

gravitational force

frictional force

electrostatic force
intermolecular force

Q83 A force which acts on a particle in such a way that there is interconversion between kinetic and potential energies is said to be _____

centripetal
inertial
conservative
kinematic

Q84 What is the torque produced by a force of $2\vec{i} - 3\vec{j} + \vec{k}$ N on a particle at radial distance of $4\vec{i} + \vec{j} - 2\vec{k}$ m from the origin a three dimensional coordinate system?

$12\vec{i} - 8\vec{j} + 7\vec{k}$ Nm
 $5\vec{i} + 3\vec{j} - 4\vec{k}$ Nm
 $\vec{i} + 8 - 2\vec{j} - \vec{k}$ Nm
 $5\vec{i} + 8\vec{j} + 14\vec{k}$ Nm

Q85 The work done by the force $2\vec{i} - 3\vec{j} + \vec{k}$ N to move a particle through a displacement of $4\vec{i} + \vec{j} - 2\vec{k}$ m is ----- J.

2
4
3
7

Q86 The angle between the vectors $6\vec{i} + 5\vec{j} + 4\vec{k}$ and $\vec{i} + 2\vec{j} + 3\vec{k}$ is

31.5°
 60.0°
 30.0°
 42.5°

Q87 For what values of q are the two vectors $\vec{i} + \vec{j} + q\vec{k}$ and $\vec{b} = q\vec{i} - 2\vec{j} + 2q\vec{k}$ parallel to each other?

0,1
0,-2
1,1
1,-2

Q88 Given that $\vec{a} = 5\vec{i} - 2\vec{j} + 3\vec{k}$;
 $\vec{b} = 3\vec{i} + \vec{j} - 2\vec{k}$; $\vec{c} = \vec{i} - 3\vec{j} + 4\vec{k}$;
evaluate the vector triple product of $\vec{a}$, $\vec{b}$ and $\vec{c}$.

$109\vec{i} + 7\vec{j} - 22\vec{k}$
 $13\vec{i} + 7\vec{j}$
 $17\vec{i} - \vec{k}$
 $121\vec{i} - 12\vec{j} + 5\vec{k}$

Q89 In spherical coordinates, which of the following relations is NOT correct?

$x = r \cos \theta \cos \phi$

$$\begin{aligned}x &= r \sin \theta \cos \phi \\ y &= r \sin \theta \sin \phi \\ z &= r \cos \theta\end{aligned}$$

Q90 Given that $\vec{a} = 5\vec{i} - 2\vec{j} + 3\vec{k}$; $\vec{b} = 3\vec{i} + \vec{j} - 2\vec{k}$; $\vec{c} = \vec{i} - 3\vec{j} + 4\vec{k}$; evaluate the scalar triple product of $\vec{a}$, $\vec{b}$ and $\vec{c}$.

- 12
-16
-12
16

Q91 If a vector function is given as $\vec{F} = xy^2\vec{i} - 2yz\vec{j} + xyz\vec{k}$, $\nabla \cdot \vec{F}$ at the point P(1,-1,2) from the origin of $\vec{F}$

$$\begin{aligned}&2(\vec{j} + \vec{k}) \\ &3\vec{j} - 4\vec{k} \\ &4\vec{j} + 3\vec{k} \\ &\vec{j} + \vec{k}\end{aligned}$$

Q92 Which of these is the correct set up of the Lagrangian for a simple pendulum?

$$\begin{aligned}L(\theta, \dot{\theta}) &= \frac{1}{2}ml^2\dot{\theta}^2 + mgl\cos\theta \\ L(\theta, \dot{\theta}) &= \frac{1}{2}ml^2\dot{\theta}^2 + mgl\sin\theta \\ L(\theta, \dot{\theta}) &= \frac{1}{2}ml^2\dot{\theta}^2 + mgl\cos\theta \\ L(\dot{\theta}, \ddot{\theta}) &= \frac{1}{2}ml^2\ddot{\theta}^2 - mgl\cos\theta\end{aligned}$$

Q93 Given that the hamiltonian function for the motion of a one-dimensional harmonic oscillator is $H = \frac{1}{2}mp^2 + \frac{1}{2}kx^2$, which of the following is the correct set of hamiltonian equation of motion for the system?

$$\begin{aligned}\frac{p}{m} &= \dot{x}, \quad kx = -\dot{p} \\ \frac{p}{m} &= \dot{x}, \quad kx = -\dot{p} \\ \frac{x}{m} &= \dot{p}, \quad kp = -\dot{x} \\ \frac{x}{m} &= \dot{p}, \quad kx = -\ddot{x}\end{aligned}$$

Q94 Which of the following is the correct Hamiltonian function for the motion of a particle in a central field?

$$\begin{aligned}H &= \frac{1}{2}\left(p_\theta^2 + \frac{p_r^2}{r^2}\right) + V(r) \\ H &= \frac{1}{2}\left(p_\theta^2 + \frac{p_r^2}{r^2}\right) + V(r) \\ H &= \frac{1}{2}\left(p_\theta^2 - \frac{p_r^2}{r^2}\right) + V(r) \\ H &= \frac{1}{2}\left(p_r^2 + \frac{p_\theta^2}{r^2}\right) + V(r)\end{aligned}$$

Q95 If a scalar field function is given as $\phi = x^2y^2 + x^3yz - yz^2$, determine $\nabla\phi$ at point P(1,-1,2) from the origin of ϕ .

$$\begin{aligned}&3\vec{i} - 4\vec{j} + 3\vec{k} \\ &-4\vec{i} + 4\vec{j} - 3\vec{k}\end{aligned}$$

$$-4\vec{i}-3\vec{j}+4\vec{k}$$

$$-4\vec{i}-4\vec{j}+3\vec{k}$$

Q96 The physical significance of the divergence of a vector is the
 measure of the outward flow of the vector from its source
 rotational capacity of the vector about its source
 perpendicularity of the vector to its direction of motion
 angular sweep the vector makes about its origin

Q97 Given that $\vec{C} = \vec{A} - \vec{B}$, what is $\vec{C} \cdot \vec{C}$?
 $A^2 + B^2 - 2AB\cos\theta$
 $A^2 - B^2 + 2AB\sin\theta$
 $A - B + AB\cos\theta$
 $+B - AB\sin\theta$

Q98 Which of the following is correct

$$\hat{i} \cdot \hat{j} = 1$$

$$\hat{i} \cdot \hat{i} = 1$$

$$\hat{i} \times \hat{j} = -\hat{k}$$

$$\hat{k} \times \hat{j} = \hat{i}$$

Q99 The dot or scalar product of a force and a displacement vectors defines

work
 power
 moment
 momentum

Q100 The central force problem is important in explaining the motion of

gases
 high energy particles
 celestial bodies
 nucleons

Q101 A vector is completely defined by _____ quantities, each with its appropriate units

Q102 $\frac{\partial H}{\partial p_k} = \dot{q}_k$, $\frac{\partial H}{\partial q_k} = -\dot{p}_k$ and $\frac{\partial H}{\partial t} = -\frac{\partial L}{\partial t}$ are known as the _____ equations of Hamilton

Q103 The equation $H = \sum_{k=1}^n p_k \dot{q}_k - L(q, \dot{q}, t)$ gives the classical Hamiltonian function for a system of particles, each of mass m_k and described by the generalized coordinates q_k . For a system of particles in a conservative field, the Hamiltonian function given represents _____ of the system

Q104 $\frac{d}{dt} \left(\frac{\partial L}{\partial \dot{q}_k} \right) = \frac{\partial L}{\partial q_k}$ represent the _____'s equations of motion for a conservative system subject to, at worst, only holonomic constraints

Q105 In the application of the variational principle to a physical system, the quantity L in the equation $L = T - V$ is called the

Q106 Given the $Q_k = F_i \frac{\partial x_i}{\partial q_k}$, Q_k represents

Q107 $p_k = \frac{\partial T}{\partial \dot{q}_k}$ represents momentum

Q108 Restrictions imposed on the free motion of a particle (or a system of particles) are generally called

Q109 The degree of is important in the design of a good suspension system of a car to ensure comfortable ride.

Q110 In a series LC circuit, the quantity $\frac{1}{LC}$ is equal to the square of the (where L and C are the inductance and capacitance respectively)

Q111 The equation of a simple pendulum is given as $m\ddot{x} + mg\frac{x}{l} = 0$. The quantity $\frac{g}{l}$ is equal to the of the

Q112 If the displacement of a simple harmonic oscillator as a function of time t given as $x = A \cos(\omega t + \phi)$, the quantity ϕ is called the constant

Q113 $\frac{1}{2} m \omega^2 A^2 \sin^2 \phi =$. Note ϕ is the potential in a conservative force field.

Q114 for small displacements from equilibrium position, the restoring force of a simple harmonic oscillator obeys law

Q115 (1) The paths of planets about the sun are elliptical in shape, with the center of the sun being located at one focus. (2) An imaginary line drawn from the center of the sun to the center of the planet will sweep out equal areas in equal intervals of time. (3) The ratio of the squares of the periods of any two planets is equal to the ratio of the cubes of their average distances from the sun. The three statements constitute what is known as 's laws

Q116 The motion of a particle in a central force field always takes place in a

Q117 In the equation $\vec{N} = \vec{r} \times \vec{F}$, $\vec{N}$ represents

Q118 For a particle moving under the central conservative force, the equation $\frac{1}{2} \dot{r}^2 + \frac{L^2}{2mr^2} + V(r) = E$ is called the energy equation

Q119 In the equation $\vec{L} = \vec{r} \times \vec{p}$, L represents

Q120 The equations
$$\dot{x}_1 = \sum_{k=1}^{3N} \frac{\partial x_1}{\partial q_k} \dot{q}_k + \frac{\partial x_1}{\partial t}$$

$$\vdots$$

$$\dot{x}_N = \sum_{k=1}^{3N} \frac{\partial x_N}{\partial q_k} \dot{q}_k + \frac{\partial x_N}{\partial t}$$
 represent the cartesian velocity components expressed in terms of general

Q121 If a system of particles is described by a set of generalised coordinates $q_1, \dots, q_{3N}$, then $\dot{q}_k$ for $k=1, 2, \dots, 3N$ is called the generalized associated with the coordinates

Q122 The generalised force associated with angular quantities is a

Q123 A particle that moves along a straight line has degree of freedom

Q124 Constraints that can be represented as functions of space and time are said to be

Q125 The mathematical term used for describing the circulation of a vector is the

Q126 If $\vec{a} \times \vec{b} = 0$, then, $\vec{a}$ and $\vec{b}$ are

Q127 The measure of the outward flow of the vector from its source is the of the vector

Q128 The dot or scalar product of a force and a displacement vectors defines

Q129 A force which acts on a particle in such a way that there is interconversion between kinetic and potential energies is said to be

Q130 If the addition of vector by geometric method gives a close figure, the the resultant of the vectors is

Q131 If $\vec{F}$ is a force and $\vec{r}$ the radius vector from the origin of the position of a particle acted upon the force, then $\vec{F} \times \vec{r}$ represents the of the force about the origin

Q132 The gradient of a scalar field function is a quantity

Q133 If the curl of a vector is zero, then the vector is said to be

Q134 Undisturbed orbital motion under the influence of a central force satisfies ___'s law of areas

Q135 The term $r \ddot{\theta}$ in the expression $\ddot{r} - r \dot{\theta}^2$ represents acceleration

Q136 The aspect of mechanics that describes the motion of particles without that regard to the dynamical laws that determine which motion actually occurs is referred to as

Q137 If $\oint \vec{F} \cdot d\vec{r} = 0$, then $\vec{F}$ is a force

Q138 The statement $T + V = E(\text{constant})$, where T is the kinetic energy and V the potential energy is known as the theorem

Q139 If $\nabla \times \vec{F} = 0$, then $\vec{F}$ is

Q140 If the work done in moving a particle from place to place in a force field is independent of the path followed by the particle, then the force field is said to be

Q141 A damped oscillation driven by external periodic impulses whose frequency is the same as the natural frequency of the oscillator

Q142 If the resistance to a simple harmonic oscillation is very strong, the motion is said to be heavily

Q143 In reality, the amplitude of a simple harmonic oscillator gradually decreases in the presence of dissipative forces. Such motion is said to be

Q144 In an LC circuit, the quantity $\frac{1}{\sqrt{LC}}$ represents

Q145 The displacement of a simple harmonic oscillator is given as $x = A \cos(\omega t + \phi)$. The quantity ω is the of the oscillation. (symbols have their usual meaning)

Q146 The term normal mode frequency is associated with oscillation

Q147 The inertial factor in the spring mass system is m . The inertial factor in the LC circuit is

Q148 Any motion that repeats itself at regular intervals of time is said to be

Q149 The equation $\frac{d}{dt} \left(\frac{1}{2} m \dot{\theta}^2 + U \right) = \frac{d}{dt} (E - V)$ is the statement of conservation of

Q150 For an oscillating system, if the restoring force obeys Hooke's law for small displacement from equilibrium position, the motion is

Q151 A damped oscillation driven by external periodic impulses whose frequency is the same as the natural frequency of the oscillator is said to -----
be heavily damped
free

resonate
be critically damped

Q152 The equation of motion of a simple harmonic oscillator is $\frac{d^2x}{dt^2} + 16x = 0$. Solve this equation subject to the initial conditions that at $t = 0$, $x = 4$ cm and $\frac{dx}{dt} = 0$.

$$x(t) = 3\cos 2t$$

$$x(t) = 4\cos 4t$$

$$x(t) = 2\cos t$$

$$x(t) = \sqrt{2}\cos\left(2t - \frac{\pi}{3}\right)$$

Q153 Which of the following is NOT a valid solution of the equation of a simple harmonic oscillator? The symbols have their usual meaning.

$$x = a\cos(\omega t + \phi)$$

$$x = a\sin(\omega t - \phi)$$

$$x = a\cos(\omega t - \phi)$$

$$x = a\sin(\omega t - \phi)$$

Q154 The equation of a simple harmonic motion is given as $\frac{d^2x}{dt^2} + \omega^2 x = 0$ where the symbols have their usual meaning. The dimension of the quantity ω^2 is

$$L^{-1}$$

$$M$$

$$T^{-2}$$

$$LT^{-2}$$

Q155 Write down the correct relation for the component of the acceleration in the $\hat{r}$ direction in the plane polar coordinate system

$$r\ddot{\theta} + 2\dot{r}\dot{\theta}$$

$$\ddot{r} + r\dot{\theta}^2$$

$$r\ddot{\theta} - 2\dot{r}\dot{\theta}$$

$$\ddot{r} - r\dot{\theta}^2$$

Q156 In the plane polar coordinate system, which of the following correctly gives the component of acceleration in the $\hat{\theta}$ direction?

$$r\ddot{\theta} + 2\dot{r}\dot{\theta}$$

$$\ddot{r} + r\dot{\theta}^2$$

$$r\ddot{\theta} - 2\dot{r}\dot{\theta}$$

$$\ddot{r} - r\dot{\theta}^2$$

Q157 In the plane polar coordinate system, which of the following correctly gives the component of velocity in the $\hat{\theta}$ direction?

$$\dot{r}$$

$$\dot{\theta}$$

$$r\dot{\theta}$$

$$r\dot{r}$$

Q158 if $\hat{r}$, $\hat{\theta}$ are the functions of θ in the plane polar coordinate system, which of the following is Not a correct relation among

θ , $\hat{r}$, and $\hat{\theta}$?

$$\hat{r} = \hat{x} \cos \theta - \hat{y} \sin \theta$$

$$\hat{r} = \hat{x} \cos \theta + \hat{y} \sin \theta$$

$$\hat{\theta} = -\hat{x} \sin \theta + \hat{y} \cos \theta$$

$$\hat{\theta} = \hat{y} \cos \theta - \hat{x} \sin \theta$$

Q159 In the plane polar coordinates, which of the following relations is NOT correct?

$$x = r \cos \theta$$

$$\theta = \sin^{-1} \frac{y}{\sqrt{x^2 + y^2}}$$

$$\theta = \cos^{-1} \frac{x}{\sqrt{x^2 + y^2}}$$

$$\theta = r \sin x$$

Q160 A particle is subjected to a force $F = (2xy + z^2) \hat{i} + x \hat{j} + 2xz \hat{k}$ in a conservative field. Find the potential function.

$$2xyz$$

$$-\frac{1}{2} (x^2 y + z^2 x)$$

$$2x^2 y z^2$$

$$-\frac{1}{2} (xy^2 + zx^2)$$

Q161 Given that the hamiltonian function for the motion of a one-dimensional harmonic oscillator is $H = \frac{1}{2m} p^2 + \frac{k}{2} x^2$, which of the following is the correct set of hamiltonian equation of motion for the system?

$$\frac{dp}{dt} = \dot{x}, \quad kx = -\dot{p}$$

$$\frac{dp}{dt} = \dot{x}, \quad kx = -\dot{p}$$

$$\frac{dx}{dt} = \dot{p}, \quad kp = -\dot{x}$$

$$\frac{dx}{dt} = \dot{p}, \quad kx = -\dot{x}$$

Q162 Which of the following is the correct Hamiltonian function for the motion of a particle in a central field?

$$H = \frac{1}{2} \left(p_{\theta}^2 + \frac{p_r^2}{r^2} \right) + V(\theta)$$

$$H = \frac{1}{2} \left(p_{\theta}^2 + \frac{p_r^2}{r^2} \right) + V(r)$$

$$H = \frac{1}{2} \left(p_{\theta}^2 - \frac{p_r^2}{r^2} \right) + V(r)$$

$$H =$$

$$H = \frac{1}{2} \left(p_r^2 + \frac{p_{\theta}^2}{r^2} \right) + V(r)$$

Q163 Which of these is the correct set up of the Lagrangian for a simple pendulum?

$$L(\theta, \dot{\theta}) = \frac{1}{2} m l^2 \dot{\theta}^2 + m g l \cos \theta$$

$$L(\theta, \dot{\theta}) = \frac{1}{2} m l^2 \dot{\theta}^2 + m g l \sin \theta$$

$$L(\theta, \dot{\theta}) = \frac{1}{2} m l^2 \dot{\theta}^2 + m g l \cos \theta$$

$$L(\dot{\theta}, \ddot{\theta}) = \frac{1}{2} m l^2 \ddot{\theta}^2 - m g l \cos \theta$$

Q164 Which of the following is NOT a correct canonical equation of Hamilton? (Symbols have their usual meaning)

$$\frac{\partial H}{\partial p_{\alpha}} = \dot{p}_{\alpha}$$

$$\frac{\partial H}{\partial p_{\alpha}} = \dot{q}_{\alpha}$$

$$\frac{\partial H}{\partial q_{\alpha}} = -\dot{p}_{\alpha}$$

$$\frac{\partial H}{\partial t} = -\frac{\partial L}{\partial t}$$

Q165 The equations $\begin{matrix} \dot{x}_1 = \sum_{k=1}^{3N} \frac{\partial x_1}{\partial q_k} \dot{q}_k + \frac{\partial x_1}{\partial t} \\ \vdots \\ \dot{x}_N = \sum_{k=1}^{3N} \frac{\partial x_N}{\partial q_k} \dot{q}_k + \frac{\partial x_N}{\partial t} \end{matrix}$

represent the cartesian velocity components expressed in terms of

- generalised coordinates and velocities
- generalised forces and acceleration
- generalised forces and torques
- generalised forces and momenta

Q166 If a system of particles is described by a set of generalised coordinates $q_1, \dots, q_{3N}$, then $\dot{q}_k$ for $k=1, 2, \dots, 3N$ is called the -----
- associated with the coordinates

- specific torque
- generalised velocity
- angular acceleration
- generalised force

Q167 Which of these is an example of a system of particles subject to constraints?

- a single particle
- 3N free particles
- rigid object
- N free particles

Q168 The generalised force associated with angular quantities is

- potential
- energy
- power
- a torque

Q169 In spherical coordinates, which of the following relations is NOT correct?

$$x = r \cos \theta \cos \phi$$

$$x = r \sin \theta \cos \phi$$

$$y = r \sin \theta \sin \phi$$

$$z = r \cos \theta$$

Q170 Two masses m_1 and m_2 connected by a spring are sliding on a frictionless plane. How many degrees of freedom has the system?

- 2
- 4
- 1
- 3

Q171 With polar coordinates (r, ϕ) as generalised coordinates,

the generalised velocities are $\left(\dot{r}, \dot{\phi} \right)$, while the velocity vector is -----

$$\begin{aligned} & \ddot{r}\hat{r} + 2\dot{r}\dot{\phi}\hat{\phi} \\ & \dot{r}\hat{r} + \dot{\phi}\hat{\phi} \\ & \dot{r}\hat{r} - \dot{\phi}\hat{\phi} \\ & \ddot{r}\hat{r} - 2\dot{r}\dot{\phi}\hat{\phi} \end{aligned}$$

Q172 How many degrees of freedom has a particle that moves along a straight line?

- 3
- 2
- 1
- 0

Q173 Constraints that can be represented as functions of space and time are said to be -----

- holonomic
- orthogonal
- inertial
- invariant

Q174 A set of coordinate axes with respect to which the position and orientation of an object can be measured is called the -----

- constraint
- degree of freedom
- orthogonality
- frame of reference

Q175 Consider two capacitively coupled circuits shown in the figure. How many normal modes are we supposed to have in the system?

- 1
- 2
- 3
- 0

Q176 The equation of motion of a simple pendulum is given as $\ddot{\theta} + \omega^2\theta = 0$. The equation of motion of a simple pendulum is given as $\ddot{\theta} + \omega^2\theta = 0$. $\omega^2 =$ -----

$$\begin{aligned} & g \\ & \sqrt{\frac{g}{l}} \\ & \frac{g}{l} \\ & \sqrt{\frac{l}{g}} \end{aligned}$$

Q177 The differential equation for the orbit of motion in an inverse square law force field is

$$\begin{aligned} & \frac{du}{d\theta} + u = \frac{k}{h^2} \\ & \frac{d^2u}{d\theta^2} + u = \frac{k}{h} \\ & \frac{d^2u}{d\theta^2} + u = 0 \end{aligned}$$

$$\frac{d^2u}{d\theta^2} + u = \frac{k}{mh^2}$$

Q178 The equation of motion of a damped oscillator is given as $m\ddot{x} + \gamma\dot{x} + kx = 0$. γ is called the damping coefficient. Which of the following is the correct unit of the damping coefficient?

- kg s^{-1}
- ms^{-1}
- kg m^{-1}
- s^{-2}

Q179 All bodies at the same place in a gravitational field experience

- same acceleration
- different acceleration
- zero acceleration
- twice the acceleration as that on the surface of the source of the field

Q180 Which of the following is a correct example of a coupled oscillator?

- an oscillating torsional pendulum
- oscillating atoms in a solid
- an acoustic oscillator
- an oscillating simple pendulum

Q181 Given that $\vec{F} = f(r)\vec{r}$, where $\vec{F}$ is the central force, $f(r)\vec{r}$ is a function of the radial distance from the origin and $\vec{r}$ is the unit vector directed radially outward from the origin, which of the following is correct?

- $f(r) < 0$ implies that the $\vec{F}$ is repulsive
- $f(r) > 0$ implies that the $\vec{F}$ is attractive
- $\vec{r} \times \vec{F} < 0$
- $\vec{r} \times \vec{F} = 0$

Q182 Which of the following is NOT a central force?

- gravitational force
- frictional force
- electrostatic force
- intermolecular force

Q183 A force which acts on a particle in such a way that there is interconversion between kinetic and potential energies is said to be _____

- centripetal
- inertial
- conservative
- kinematic

Q184 What is the torque produced by a force of $2\vec{i} - 3\vec{j} + \vec{k}$ N on a particle at radial distance of $4\vec{i} + \vec{j} - 2\vec{k}$ m from the origin a three dimensional coordinate system?

- $12\vec{i} - 8\vec{j} + 7\vec{k}$ Nm
- $5\vec{i} + 3\vec{j} - 4\vec{k}$ Nm

$$(\vec{i} + 8 - 2\vec{j} - \vec{k}) \cdot \vec{k} \text{ Nm}$$

$$5(\vec{i} + 8\vec{j} + 14\vec{k}) \cdot \vec{k} \text{ Nm}$$

Q185 The work done by the force $2\vec{i} - 3\vec{j} + \vec{k}$ N to move a particle through a displacement of $4\vec{i} + \vec{j} - 2\vec{k}$ m is ----- J.

- 2
4
3
7

Q186 The angle between the vectors $6\vec{i} + 5\vec{j} + 4\vec{k}$ and $\vec{i} + 2\vec{j} + 3\vec{k}$ is

$$31.5^\circ$$

$$60.0^\circ$$

$$30.0^\circ$$

$$42.5^\circ$$

Q187 For what values of q are the two vectors $\vec{i} + \vec{j} + q\vec{k}$ and $\vec{b} = q\vec{i} - 2\vec{j} + 2q\vec{k}$ parallel to each other?

- 0,1
0,-2
1,1
1,-2

Q188 Given that $\vec{a} = 5\vec{i} - 2\vec{j} + 3\vec{k}$; $\vec{b} = 3\vec{i} + \vec{j} - 2\vec{k}$; $\vec{c} = \vec{i} - 3\vec{j} + 4\vec{k}$; evaluate the vector triple product of $\vec{a}$, $\vec{b}$ and $\vec{c}$.

$$109\vec{i} + 7\vec{j} - 22\vec{k}$$

$$13\vec{i} + 7\vec{j}$$

$$17\vec{i} - \vec{k}$$

$$121\vec{i} - 12\vec{j} + 5\vec{k}$$

Q189 In spherical coordinates, which of the following relations is NOT correct?

$$x = r \cos \theta \cos \phi$$

$$x = r \sin \theta \cos \phi$$

$$y = r \sin \theta \sin \phi$$

$$z = r \cos \theta$$

Q190 Given that $\vec{a} = 5\vec{i} - 2\vec{j} + 3\vec{k}$; $\vec{b} = 3\vec{i} + \vec{j} - 2\vec{k}$; $\vec{c} = \vec{i} - 3\vec{j} + 4\vec{k}$; evaluate the scalar triple product of $\vec{a}$, $\vec{b}$ and $\vec{c}$.

- 12
-16
-12
16

Q191 If a vector function is given as $\vec{F} = xy^2\vec{i} - 2yz\vec{j} + xyz\vec{k}$, $\nabla \cdot \vec{F}$ at the point P(1,-1,2) from the origin of $\vec{F}$

$$\begin{aligned} & \$2(\vec{J}+\vec{K})\$ \\ & \$3\vec{J}-4\vec{K}\$ \\ & \$4\vec{J}+3\vec{K}\$ \\ & \$\vec{J}+\vec{k}\$ \end{aligned}$$

Q192 Which of these is the correct set up of the Lagrangian for a simple pendulum?

$$\begin{aligned} & \$L\left(\theta, \ddot{\theta}\right)=\frac{1}{2} m l^2 \ddot{\theta}^2+m g l \cos \theta\$ \\ & \$L\left(\theta, \dot{\theta}\right)=\frac{1}{2} m l^2 \ddot{\theta}^2+m g l \sin \theta\$ \\ & \$L\left(\theta, \dot{\theta}\right)=\frac{1}{2} m l^2 \dot{\theta}^2+m g l \cos \theta\$ \\ & \$L\left(\dot{\theta}, \ddot{\theta}\right)=\frac{1}{2} m l^2 \ddot{\theta}^2-m g l \cos \theta\$ \end{aligned}$$

Q193 Given that the hamiltonian function for the motion of a one-dimensional harmonic oscillator is $H=\frac{1}{2} m p^2+\frac{k}{2} x^2$, which of the following is the correct set of hamiltonian equation of motion for the system?

$$\begin{aligned} & \$\frac{p}{m}=\ddot{x}, \$kx=-\ddot{p}\$ \\ & \$\frac{p}{m}=\dot{x}, \$kx=-\dot{p}\$ \\ & \$\frac{x}{m}=\dot{p}, \$kp=-\dot{x}\$ \\ & \$\frac{x}{m}=\dot{p}, \$kx=-\ddot{x}\$ \end{aligned}$$

Q194 Which of the following is the correct Hamiltonian function for the motion of a particle in a central field?

$$\begin{aligned} & \$H=\frac{1}{2}\left(p_{\theta}^2+\frac{p_r^2}{\theta^2}\right)+V(\theta)\$ \\ & \$H=\frac{1}{2}\left(p_{\theta}^2+\frac{p_r^2}{\theta^2}\right)+V(r)\$ \\ & \$H=\frac{1}{2}\left(p_{\theta}^2+\frac{p_r^2}{\theta^2}\right)+V(r)\$ \\ & \$H=\frac{1}{2}\left(p_r^2+\frac{p_{\theta}^2}{r^2}\right)+V(r)\$ \end{aligned}$$

Q195 If a scalar field function is given as $\phi=x^2 y^2+x^3 y z-y z^2$, determine $\nabla \phi$ at point P(1,-1,2) from the origin of ϕ .

$$\begin{aligned} & \$3\vec{i}-4\vec{j}+3\vec{k}\$ \\ & \$-4\vec{i}+4\vec{j}-3\vec{k}\$ \\ & \$-4\vec{i}-3\vec{j}+4\vec{k}\$ \\ & \$-4\vec{i}-4\vec{j}+3\vec{k}\$ \end{aligned}$$

Q196 The physical significance of the divergence of a vector is the
 measure of the outward flow of the vector from its source
 rotational capacity of the vector about its source
 perpendicularity of the vector to its direction of motion
 angular sweep the vector makes about its origin

Q197 Given that $\vec{C}=\vec{A} \times \vec{B}$, what is $\vec{C} \cdot \vec{C}$?

$$\begin{aligned} & \$A^2+B^2-2 A B \cos \theta\$ \\ & \$A^2-B^2+2 A B \sin \theta\$ \\ & \$A \cdot B+A B \cos \theta\$ \\ & \$+B \cdot A B \sin \theta\$ \end{aligned}$$

Q198 Which of the following is correct

$$\hat{i} \cdot \hat{j} = 1$$

$$\hat{i} \cdot \hat{i} = 1$$

$$\hat{i} \times \hat{j} = \hat{k}$$

$$\hat{k} \times \hat{j} = \hat{i}$$

Q199 The dot or scalar product of a force and a displacement vectors defines

work

power

moment

momentum

Q200 The central force problem is important in explaining the motion of

gases

high energy particles

celestial bodies

nucleons

noungeeks.com