## Join group: T.me/NOUNSTUDENTSFORUM CLICK TO DOWNLOAD MORE TMA PQ

conjugated

[CHM318] β-Carotene was first isolated from carrots by a Scientist Wackarodder in the yearââ,¬Â¹. 1837
[CHM318] Gonane is the simple possible structure ofââ,¬Â¦Ã¢â,¬Â¦ steroids
[CHM318] The terpenoids called carotenoids hascarbon atoms 40
[CHM318] The structure of camphor was established inââ,¬Â¦. 1893
[CHM318] The other name for isoprenoids is Sterculine
[CHM318] The molecules whose structure is based on isoprene units is called $\hat{A}$ ¢ $\hat{a}$ , ¬ $\hat{A}$ ¦ $\hat{A}$ ¢ $\hat{a}$ , ¬ $\hat{A}$ ! Terpenes
[CHM318] The diterpenoids has a total ofcarbon atoms 20
[CHM318] The Mevalonic acid-independent pathway was discovered in $\tilde{A} \not c \hat{a}, \neg \hat{A} \mid .$ 1980s
[CHM318] Isoprene unit hascarbon atoms 5

Whatsapp: 08089722160 or click here for TMA assistance

Practice E-exams & Chat with course mates on noungeeks.net