

determination of urinary flow rate over time it is a simple way to measure bladder emptying.
Uroflowmetry

[NSC402] The type of injection that is used to deliver drugs for optic neuritis, papillitis and posterior uveitis is called *retrobulbar*.

[NSC402] The kidneys can maintain a constant filtration rate changes in arterial pressure from *80 to 180 mmHg* to produce little change in blood flow and filtration rate in glomerulus.

[NSC402] In urine formation, *0.8* of materials that are reabsorbed take place in proximal convoluted tubules cells lining

[NSC402] New born glaucoma that is also called true congenital glaucoma accounts for *0.25* of all cases of glaucoma

[NSC402] *Keratoconus* is a non-inflammatory bilateral form of corneal degeneration in which the cornea becomes thin and protrudes anteriorly.

[NSC402] *Amethocaine* is an example of topical anaesthetic agents are useful for examination and treatment of simple procedures such as removal of foreign bodies from the cornea.

[NSC402] Examples of antiviral drug that is effective against the herpes simplex keratitis is *Trifluorothymidine*

[NSC402] Examining the eyelids can help to discover the following abnormalities except cataract

[NSC402] Retina disorder that is characterized by multiple superficial retinal haemorrhages, involving posterior part of the fundus is called *Subacute retinitis of Roth*

Whatsapp: 08089722160 or click here for TMA assistance

Practice E-exams & Chat with course mates on [noungeeks.net](https://www.noungeeks.net)