

Korsakoff Psychosis

[NSC503] This is a behavioral pattern of drug use characterized by overwhelming involvement with the use of a drug.

Addiction

[NSC503] It is characterized by mental confusion, tremor, sensory hyperactivity, visual hallucinations.

Delirium Tremens

[NSC503] The bodily metabolism has become adapted to the presence of alcohol.

Alcohol Addiction

[NSC503] _____ implies overzealous or indiscreet administration of drugs by physicians.

Misuse

[NSC503] _____ is characterized by a pattern of pathologic use lasting for at least an month and causing impairment in social and occupational functioning.

Abuse

[NSC503] There is disorientation, associated with paralysis of ocular muscles, nystagmus, ataxia. It is due to acute deficiency of vitamin B1.

Wernicke's Encephalopathy

[NSC503] It is a compulsive drug-seeking behaviour in which the individual uses the drug repetitively for personal satisfaction.

Dependence

[NSC503] There is a psychological dependence on alcohol which is needed to relieve physical or emotional discomfort.

Habitual Symptomatic Excessive Drinkers

[NSC503] When an individual in crisis perceived threat acts as a precipitant that generates increased anxiety.

Phase I

Whatsapp: 08089722160 or click here for TMA assistance

Practice E-exams & Chat with course mates on noungeeks.net