

the number of neutrons and protons

[PHY402] An element X has two isotopes, A_1X and ${}^{A+2}_2X$ present in the ratio 3:1. Calculate the relative atomic mass of X is 21.5

[PHY402] The ability of the γ ray to be able to dislodge electrons from both the outer and inner orbit are done through the following ways except resistance effect

[PHY402] The amount of energy released when 0.5kg of uranium is burnt completely. Take $c = 3 \times 10^8$ ms⁻¹
 4.5×10^{11} J

[PHY402] An element X has 127 neutrons and 82 electrons. Its atomic number is 82

[PHY402] The half life radioactive nuclide is 20 years. Calculate the fraction that remains after ten years.
0.3465

[PHY402] When atoms undergoes a beta decay, the atomic number of the nucleus increase by one

[PHY402] The decay constant of a certain radioactive nuclide is 0.0200m⁻¹. Calculate the fraction that would remain from the sample of 0.2kg of the substance after 3yrs
0.95

[PHY402] The mass number of an atom is the sum of number of protons and neutrons in the nucleus of the atom

[PHY402] The half-life of a radioactive nuclide is 28 years. Calculate the decay constant.
 2.475×10^{-12} s⁻¹

Whatsapp: 08089722160 or click here for TMA assistance

Practice E-exams & Chat with course mates on noungeeks.net